

## Photochemistry of Metalloporphyrin Carbene Complexes

Christopher J. Ziegler and Kenneth S. Suslick\*

School of Chemical Sciences  
University of Illinois at Urbana-Champaign  
Urbana, Illinois 61801

Received December 6, 1995

The photochemistry of metalloporphyrins and other tetrapyrrolic systems continues to attract considerable interest.<sup>1</sup> There have been numerous studies on the photoactivity of various porphyrin complexes with single-bonded axial ligands, including halogens, oxo-anion species, molecular oxygen,<sup>2</sup> and a variety of metals, including chromium, manganese, and iron.<sup>3</sup> The photochemistry of metalloporphyrins with multiple-bonded axial ligands has not been explored. We report here the photochemistry of several iron porphyrin carbene and vinylidene complexes. Irradiation of these complexes with visible light cleaves the iron–carbon double bond, resulting in a four-coordinate iron(II) porphyrin and a free carbene. This chemistry is unique in the area of transition metal carbene or alkylidene compounds.

Porphyrins with multiple-bonded axial ligands are especially relevant to both biological and catalytic oxidative processes.<sup>4</sup> Iron porphyrin carbene and nitrene species are isoelectronic with a terminal iron–oxo complex, the proposed intermediate of the cytochrome P-450 monooxygenases and its synthetic analogs. Most recently, the catalytic cyclopropanation of olefins by iron porphyrins using ethyl diazoacetate was thought to involve a porphyrin carbene complex.<sup>5</sup>

We prepared Fe(TPP)CCl<sub>2</sub>, Fe(TPP)CBr<sub>2</sub>, Fe(TPP)CClF, and Fe(TPP)CC(C<sub>6</sub>H<sub>4</sub>Cl)<sub>2</sub> under anaerobic conditions following Mansuy's procedure using iron metal as the reductant (TPP = 5,10,15,20-tetraphenylporphyrinate).<sup>6</sup> These compounds were further purified via crystallization from benzene/pentane under argon in an inert atmosphere box and then fully characterized.<sup>7</sup> To examine the photochemistry of these complexes, degassed benzene solutions (~1 × 10<sup>-4</sup> M) were irradiated with a 300 W Xe arc lamp, which was filtered to remove both infrared and ultraviolet (<360 nm) light in order to prevent sample heating and porphyrin bleaching, respectively. In addition, the photolysis cells were thermostated to an internal solution temperature of 20 °C.

(1) (a) *Photochemistry of Polypyridine and Porphyrin Complexes*; Kalyanasundaram, K., Ed.; Academic: New York, 1992. (b) Blauer, G.; Sund, H. *Optical Properties and Structure of Tetrapyrroles*; W. De Gruyter: Berlin, 1985. (c) Mauzerall, D. In *The Porphyrins*; Dolphin, D., Ed.; Academic: New York, 1979; Vol. 5, p 29 ff.

(2) (a) Imamura, T.; Jin T.; Suzuki, T.; Fujimoto, M. *Chem. Lett.* **1985**, 847. (b) Hendrickson, D. N.; Kinnaird, M. G.; Suslick, K. S. *J. Am. Chem. Soc.* **1987**, 109, 1243. (c) Jin, T.; Suzuki, T.; Imamura, T.; Fujimoto, M. *Inorg. Chem.* **1987**, 26, 1280. (d) Suslick, K. S.; Watson, R. A. *Inorg. Chem.* **1991**, 30, 912. (e) Suslick, K. S.; Watson, R. A.; Wilson, S. R. *Inorg. Chem.* **1991**, 30, 2311. (f) Proniewicz, L. M.; Bajdor, K.; Nakamoto, K.; *J. Phys. Chem.* **1986**, 90, 1760.

(3) (a) Suslick, K. S.; Watson, R. A. *New J. Chem.* **1992**, 16, 633. (b) Suslick, K. S.; Acholla, F. A.; Cook, B. R. *J. Am. Chem. Soc.* **1987**, 109, 2818. (c) Buchler, J. W.; Dreher, C. *Z. Naturforsch.* **1984**, 39b, 222.

(4) Sheldon, R. A., Ed. *Metalloporphyrins in Catalytic Oxidations*; Marcel Dekker: New York, 1994.

(5) Wolf, J. R.; Hamaker, C. G.; Djukic, J-P.; Kodadek, T.; Woo, L. K. *J. Am. Chem. Soc.* **1995**, 117, 9194.

(6) (a) Mansuy, D.; Lange, M.; Chottard, J. C.; Guerin, P.; Morliere, P.; Brault, D.; Rougee, M. *J. Chem. Soc., Chem. Commun.* **1977**, 648. (b) Mansuy, D.; Lange, M.; Chottard, J. C.; Bartoli, J. F.; Chevrier, B.; Weiss, R. *Angew. Chem., Int. Ed. Engl.* **1978**, 17, 780. (c) Mansuy, D. *Pure Appl. Chem.* **1980**, 52, 681. (d) Mansuy, D.; Lange, M.; Chottard, J. C. *J. Am. Chem. Soc.* **1978**, 100, 3214.

(7) UV-Vis, <sup>1</sup>H NMR, FDMS, and elemental analysis were used to confirm the identity and purity of these species, which agreed completely with the prior literature.





Figure 1. Photolysis of Fe(TPP)CCl<sub>2</sub> in benzene under argon at 25 °C.

Table 1. Photolysis of Fe(TPP)CX<sub>2</sub> in the Presence of Alkene Substrates

Porphyrin	Substrates, Products, and Yields			
Fe(TPP)CCl <sub>2</sub>	 82%	 80%	 80%	 79%
Fe(TPP)CBr <sub>2</sub>	 59%	 59%	 64%	 55%
Fe(TPP)CClF	 90%	 89%	 85%	 86%

In all cases, irradiation of the parent compounds cleanly produced a new porphyrin identified as Fe(TPP) ( $\lambda_{\text{max}} = 419, 442, \text{ and } 537 \text{ nm}$ ), as shown in Figure 1 for Fe(TPP)CCl<sub>2</sub>. Both the Soret and Q-bands of the parent compounds are photoactive.<sup>8</sup> The formation of Fe(TPP) was confirmed by addition of pyridine after photolysis of the carbene or vinylidene complexes, yielding the bispyridine complex, Fe(TPP)(py)<sub>2</sub>. Also as expected, exposure of photolysed solutions to oxygen resulted in the formation of the  $\mu$ -oxo dimer species, [Fe(TPP)]<sub>2</sub>O.

The production of Fe(TPP) in this reaction is indicative of homolytic cleavage of the metal–carbon double bond and production of a carbene, as in the following reaction:













The transient formation of free carbenes was confirmed by addition to various substrates (Tables 1 and 2), with products monitored by capillary GC and GCMS. The reactivity of free carbenes with alkene substrates has been thoroughly investi-

(8) The quantum yields for these compounds are comparable to those of other metalloporphyrins with axial ligands. These yields upon irradiation of the Soret band ranged from  $1 \times 10^{-4}$  to  $4 \times 10^{-3}$  with  $\text{CBr}_2 < \text{CCl}_2 < \text{CC}(\text{C}_6\text{H}_4\text{Cl})_2 < \text{CClF}$ , which tracks the relative stabilities of the four resulting carbene fragments. The Q-bands, upon selective irradiation, were also found to be photoactive.

(9) (a) Kirmse, W. *Carbene Chemistry*, 2nd ed.; Academic: New York, 1971; pp 267–362. (b) Parham, W. E.; Schweizer, E. E. *Org. React.* **1963**, 13, 55.

**Table 2.** Photolysis of Fe(TPP)CX<sub>2</sub> under Competitive Conditions

Porphyrin	Substrate	Product	Photolysis Product Ratio	Base Induced Product Ratio
Fe(TPP)CCl <sub>2</sub>			0.22	0.22
Fe(TPP)CCl <sub>2</sub>			0.14	0.14
Fe(TPP)CClF		 <i>syn F vs. syn Cl products</i>	0.55	0.55
Fe(TPP)CClF		 <i>syn F vs. syn Cl products</i>	0.82	0.82
Fe(TPP)CClF		 <i>syn F vs. syn Cl products</i>	0.50	0.50

gated,<sup>9,10</sup> and the selectivity of product formation can be used as a test of the formation of free carbenes. A number of alkenes were selected to corroborate the formation of carbenes from the photolysis of the metalloporphyrin carbene complexes. Irradiation of Fe(TPP)CCl<sub>2</sub>, Fe(TPP)CBr<sub>2</sub>, or Fe(TPP)CClF in the presence of alkenes produced dihalocyclopropanes (Table 1). All of these additions occurred in good yield in dilute solution, and the relative yields from the three complexes reflect the relative reactivity of the three free carbene fragments in solution. The photolysis of more concentrated solutions of these three species resulted in the formation of Fe(TPP)X in addition to the cyclopropanes and Fe(TPP). The production of Fe(TPP)X in solution results from halocarbene attack on unreacted Fe(TPP)CX<sub>2</sub> and can be inhibited by photolysis in the presence of high alkene substrate concentration.

(10) (a) Skell, P. S.; Cholod, M. S. *J. Am. Chem. Soc.* **1969**, *91*, 6035. (b) Skell, P. S.; Cholod, M. S. *J. Am. Chem. Soc.* **1969**, *91*, 7131.

(11) (a) Cook, B. R.; Reinert, T. J.; Suslick, K. S. *J. Am. Chem. Soc.* **1986**, *108*, 7281. (b) Suslick, K. S.; Cook, B. R. *J. Chem. Soc., Chem. Commun.* **1987**, 200. (c) Suslick, K. S. In *Activation and Functionalization of Alkanes*; Hill, C. L., Ed.; John Wiley and Sons: New York, 1989; pp 219–241. (d) Collman, J. P.; Zhang, X.; Lee, V. J.; Uffelman, E. S.; Brauman, J. I. *Science* **1993**, *261*, 1404. (e) Suslick, K. S.; Van Deusen-Jeffries, S. In *Comprehensive Supramolecular Chemistry*; Lehn, J. M., Ed.; Elsevier: London, 1996; Vol. 5, in press.

(12) Hartzler, H. D. *J. Am. Chem. Soc.* **1964**, *86*, 526.

(13) (a) Pourreau, D. B.; Geoffroy, G. L. Photochemistry of Alkyl, Alkylidene, and Alkylidene Complexes of the Transition Metals. *Advances in Organometallic Chemistry*; Academic: New York, 1985; Vol. 24, pp 326–340.

(14) (a) Foley, H. C.; Strubinger, L. M.; Targos, T. S.; Geoffroy, G. L. *J. Am. Chem. Soc.* **1983**, *105*, 3064. (b) Fong, L. K.; Cooper, N. J. *J. Am. Chem. Soc.* **1984**, *106*, 2595. (c) Ofele, K.; Roos, E.; Herberhold, M. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1976**, *31B*, 1070. (d) Rieke, R. D.; Kojima, H.; Ofele, K. *J. Am. Chem. Soc.* **1976**, *98*, 6735. (e) Casey, C. P.; Shusterman, A. J.; Vollendorf, N. W.; Haller, K. J. *J. Am. Chem. Soc.* **1982**, *104*, 2417.

(15) (a) McGuire, M. A.; Hegedus, L. S. *J. Am. Chem. Soc.* **1982**, *104*, 5538. (b) Hegedus, L. S.; McGuire, M. A.; Schultze, L. M.; Yujin, C.; Anderson, O. P. *J. Am. Chem. Soc.* **1984**, *106*, 2680. (c) Borel, C.; Hegedus, L. S.; Krebs, J. Satoh, Y. *J. Am. Chem. Soc.* **1987**, *109*, 1101. (d) Hegedus, L. S.; deWeck, G.; D'Andrea, S. *J. Am. Chem. Soc.* **1988**, *110*, 2122. (e) Lastra, E.; Hegedus, L. S. *J. Am. Chem. Soc.* **1993**, *115*, 87. (f) Hegedus, L. S. *Acc. Chem. Res.* **1995**, *28*, 299.

(16) (a) Rooney, A. D.; McGarvey, J. J.; Gordon, K. C. *Organometallics* **1995**, *14*, 107. (b) McCormick, F. B.; Kiel, W. A.; Gladysz, J. A. *Organometallics* **1982**, *1*, 405.


(17) Gouterman, M. In *The Porphyrins*; Dolphin, D., Ed.; Academic: New York, 1979; Vol. 3, pp 1–165.

(18) Tatsumi, K.; Hoffmann, R. *Inorg. Chem.* **1981**, *20*, 3771.

In the photochemistry of Fe(TPP)CX<sub>2</sub>, cyclopropanation could conceivably occur either through free dihalocarbene production or through a metal-mediated carbenoid intermediate. In the first case, photoexcitation would liberate the carbene fragment, which would then go on to react with the alkene substrate. In the second possibility, the metal center would play a distinct role in the reaction, perhaps via a metallocycle intermediate. The mechanism of photochemical cyclopropanation with these complexes was determined by competitive substrate addition, from either intra- or intermolecular selectivity. Similar approaches have been found useful in mechanistic studies of epoxidations by metalloporphyrin oxo complexes.<sup>11</sup>

In the irradiation of Fe(TPP)CCl<sub>2</sub>, both the selectivity of an intramolecular competition, (*S*)-(–)-limonene (first entry in Table 2), and an intermolecular competition, cyclohexene vs 1-hexene (second entry in Table 2), were examined. In both cases, the product ratios matched exactly that of free CCl<sub>2</sub> produced from the reaction of base with CHCl<sub>3</sub> conducted at the same temperature (20 °C). In addition, photolysis of Fe(TPP)CClF with alkene substrates produced both *syn*-fluoro and *syn*-chloro products, whose ratios could also be compared to the reaction of CHCl<sub>2</sub>F with base. As before, no significant difference in product ratios was found between the metalloporphyrin complex photolysis and the free carbene reactions. This similarity in selectivity indicates that upon photoactivation the parent compound undergoes homolytic dissociation with the formation of free carbene.

Unlike the photolyses of the dihalocarbene iron porphyrins, the vinylidene complex Fe(TPP)CC(*p*-C<sub>6</sub>H<sub>4</sub>Cl)<sub>2</sub> did not liberate a reactive fragment that added to alkene substrates. Instead, upon photolysis the vinylidene complex produced di(*p*-chlorophenyl)acetylene:



The yield of the above acetylene, confirmed by GCMS, occurred in quantitative yield (99%). This phenyl migration has been seen before in a number of other organic reactions that produce the vinylidene fragment as an intermediate.<sup>12</sup>

The photochemical generation of a free carbene fragment from a transition metal carbene complex has not been observed previously.<sup>13</sup> Although the photochemistry of both Fischer and Schrock type carbene complexes has been investigated, no examples of homolytic carbene dissociation have been found. In Fischer type carbenes, photoexcitation often leads to loss of carbonyl and subsequent reaction with substrates,<sup>14</sup> or migration and insertion of a carbonyl into the metal-carbon bond, as seen in the recent chemistry of Hegedus and co-workers.<sup>15</sup> In other carbene and alkylidene compounds, such as (CO)<sub>5</sub>W=C(OR)-R' and [Cp(NO)(PPh<sub>3</sub>)Re=ChPh]<sup>+</sup>, isomerization of the metal-carbon double bond occurs.<sup>16</sup> In the case of the metalloporphyrin carbene complexes, which should be referred to as "Mansuy type" carbene complexes, the porphyrin excitation initiates the metal carbon bond cleavage. We believe that this is due to the mixing between the π\* orbitals of the porphyrin ring and the iron-carbon orbitals, which is also responsible for the *hypso* type spectrum<sup>17</sup> (i.e., blue-shifted) seen in these complexes.<sup>18</sup> *Hypso* spectra are a common trait in other photodissociative porphyrin complexes, most notably CO complexes.<sup>1</sup>

**Acknowledgment.** This work was supported by a grant from the National Institutes of Health (HL25934) and a National Science Foundation Graduate Fellowship (C.J.Z.).

JA954097E